

Inpatient Reasons for Referral to Palliative Specialist

1. Needs someone to support or lead a challenging advance care planning or goals of care discussion¹
2. Has progressive disease where uncontrolled symptoms interfere with quality of life or performance status¹
3. Has a cancer diagnosis and has failure to thrive or frailty¹
4. Patient or family requests a palliative care consult¹
5. Has a life limiting cancer and poor functional status (ECOG of 3) and the patient or health care agent has declined a hospice referral
6. Has chemotherapy-refractory advanced cancer with a good functional status and is not yet being referred for hospice¹
7. Has any stage cancer diagnosis, but quality of life and/or survival is limited by debility, has frequent hospitalizations and/or other concurrent chronic medical issues such that prognosis is less than 1 year¹
8. Has a life-limiting oncologic illness and prolonged hospital stay (greater than 7 days) without evidence of clinical improvement¹
9. Stage IV malignancy or refractory hematologic malignancy in addition to poor functional status¹
10. Current or past enrollee of hospice program¹

ICU - Reasons for Referral to Palliative Specialist

1. ICU stay longer than 7 days without evidence of improvement¹
2. Cardiac arrest (either in or out of hospital) ¹
3. Multi-system organ failure (3 or more) ¹
4. Stage IV malignancy or refractory hematologic malignancy¹
5. Poor neurologic prognosis with low chance of meaningful recovery¹
6. Inability to wean a patient from the ventilator¹
7. Team/family discussing tracheostomy, feeding tube or long term care placement¹
8. Current or past enrollee of hospice program¹
9. There is a family disagreement with the medical team, with the patient's advance directive, or with each other¹
10. Patient or family requests a palliative care consult¹

Reasons for Referral to Hospice Care

1. The patient is no longer able to come into the oncology clinic for visits due to debility, symptoms etc.²
2. Patient with chemotherapy-refractory metastatic solid tumor malignancy or refractory hematologic malignancy
3. Life expectancy less than 6 months and the patient's goals of care are focused on comfort⁴
4. Poor performance status (ECOG 2 or more) which inhibits the use of chemotherapy
5. Patient with a cancer diagnosis, other serious chronic comorbidities, debility and/or frequent hospitalizations as well as a life expectancy of less than 6 months^{2, 4}

¹Weissman DE, Meier DE. Identifying patients in need of a palliative care assessment in the hospital setting: A consensus report from the Center to Advance Palliative Care. *J Palliat Med.* 2011;14:17–23.

²National Hospice and Palliative Care Organization. http://www.nhpco.org/sites/default/files/public/NHHP_Palliative_Care_Pocket-Card.pdf. Accessed January 22, 2016.

³Prigerson, H., Bao, Y., Shah, M. et al, Chemotherapy use, performance status, and quality of life at the end of life. *JAMA Oncol.* 2015.

⁴Medicare Guidelines for Hospice eligibility.

Any clinician seeking to apply or consult the Coleman Supportive Oncology Initiative Follow Up Reference is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. The Coleman Foundation makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way.